

# The Comparison of Patient Satisfaction after “Surgery First” and Conventional Orthognathic Surgery: An Original Research

## Abstract

**Aims and Objectives:** We evaluated the factors which affect patient satisfaction and their expectations toward “surgery first” and conventional orthognathic surgery. **Materials and Methods:** Questionnaires consisting of 17 questions were given to the subjects postoperatively who had undergone Le fort 1 maxillary superior impaction for skeletal gummy smile correction with conventional orthognathic surgery and “surgery first” approach. Eleven-point rating scale based on a (visual analog scale; 0 = poor; 10 = excellent) were used to answer six questions. Ten closed-form questions were also included as well as one open question for “further remarks.” **Results:** Nineteen patients (12 females, 07 males; mean age,  $23.4 \pm 4.9$  [standard deviation] years) gave their consent to participate in the study. The intention to undergo surgery only for esthetics was noted in 42.11% of patients; only improvement of chewing function in 21.05% and both in 36.84%. **Conclusion:** The most common factor for patient satisfaction after “surgery first” and conventional orthognathic surgery was facial esthetics, but masticatory function and even psychological aspects should be considered equally when planning surgery. The patient satisfaction was more in “surgery first” than conventional orthognathic surgery. The timing of treatment and immediate results are important factors toward patient satisfaction.

**Keywords:** Esthetics, conventional orthognathic surgery, Le fort 1 osteotomy, patient satisfaction, “Surgery first”

## Introduction

The conventional approach to orthognathic surgery involves a variable length of preoperative orthodontics, orthognathic surgery, and a relatively stable period of postoperative orthodontics.<sup>[1]</sup> The importance of presurgical orthodontics rests on the positioning of the skeletal component during surgery which may be limited by inappropriate dental alignment. However, orthodontic preparation lasts 15–24 months, which involves progressive deterioration of facial esthetics and dental function and thereby causes significant patient discomfort.<sup>[2,3]</sup>

An alternative methodology is the “surgery-first” approach (SFA). It was first proposed by Nagasaka *et al.* in 2009, this method proceeds with orthognathic surgery directly without presurgical orthodontics and is followed by regular postsurgical dental corrections.<sup>[4,5]</sup> The concept implies that the most of the orthodontic treatment is performed postoperatively and compared with the

conventional approach. Thereby leading to a significant decrease in the overall total treatment time. This fact has a very positive influence on patients’ satisfaction with treatment.<sup>[6,7]</sup> The high orthodontic efficiency observed in SFA cases is due to the combination of two factors. First, the correction of the skeletal base, the complexity of orthodontic treatment is decreased, and soft tissue imbalances that might interfere with certain orthodontic tooth movements are eliminated from the beginning. Second, tooth movement is accelerated owing to the greater postoperative metabolic turnover. The optimal esthetic results, the significant reduction in overall treatment duration, and patient satisfaction led to a reasonable, cost-effective method to manage skeletal discrepancies in selected cases and that it has the potential to become a standard approach to orthognathic surgery in the future.<sup>[8,9]</sup>

Dentofacial deformities have a negative impact on many aspects of life. These include social interactions, professional

**Pritam Mohanty,  
Swati Saraswata  
Acharya<sup>1</sup>,  
Nivedita Sahoo,  
Sushila Sah,  
Sanjeeb Kumar  
Sahu<sup>1</sup>**

*Department of Orthodontics,  
Kalinga Institute of Dental  
Sciences, KIIT University,  
<sup>1</sup>Department of Orthodontics,  
Institute of Dental Sciences,  
SOA University, Bhubaneswar,  
Odisha, India*

### Address for correspondence:

*Dr. Pritam Mohanty,  
Department of Orthodontics,  
Kalinga Institute of Dental  
Sciences, KIIT University,  
Bhubaneswar - 751 024,  
Odisha, India.  
E-mail: drpritamohanty@  
gmail.com*

### Access this article online

#### Website:

www.apospublications.com

DOI: 10.4103/apos.apos\_18\_17

#### Quick Response Code:



**How to cite this article:** Mohanty P, Acharya SS, Sahoo N, Sah S, Sahu SK. The comparison of patient satisfaction after “surgery first” and conventional orthognathic surgery: An original research. APOS Trends Orthod 2017;7:123-9.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

career, love life, and personality.<sup>[10]</sup> Patients with skeletal problems may have a lower quality of life. The perception and expectations of a patient are dependent on several factors. These may be not only esthetics and function but also psychological factors.<sup>[11-13]</sup>

Hence, evaluation of different subjective and objective factors regarding patient satisfaction after conventional orthognathic surgery and SFA is needed. The comparison of patient's satisfaction after SFA and conventional orthognathic surgery is also needed to be assessed. Keeping these things in mind, a questionnaire was constructed to assess patient's satisfaction toward facial appearance after SFA and conventional orthognathic surgery.

## Materials and Methods

Nineteen patients (12 females, 7 males) agreed to participate in the study. Informed consent was obtained from all the participants, and the confidentiality of the subjects was guaranteed. The questionnaires were made by a group of orthodontists and their patients who discussed on the experience and expectations of a patient undergoing orthognathic surgery. The questionnaires were regarding different subjective and objective factors were given to the subjects postoperatively who had undergone Le Fort I maxillary superior impaction for skeletal gummy smile correction with conventional surgery and SFA. Questionnaires were explained to the patients regarding their perceptions for improvement of orthognathic surgery. The Six questions were answered using an 11-point rating scale based on a (visual analog scale; 0 = poor; 10 = excellent). Ten closed-form questions were included as well as one open question for "further remarks." [Figures 1-3].

**QUESTIONNAIRE**

Please read the following statements carefully. In order to find out how important each of the statements is to you.

1. What was your primary intention to undergo orthognathic surgery? Please mark one of the four given answers:
  - a. Aesthetic improvement only
  - b. improvement of chewing/masticatory function only
  - c. both
  - d. None/don't know
2. What type of orthognathic surgery you have undergone?
  - a) Conventional orthognathic surgery (6-8 months later)
  - b) Surgery first approach (immediate surgery). Please mark.
3. Have you given two types of surgical options before the treatment (conventional orthognathic surgery and surgery first approach)? (Yes or No)
4. Are you satisfied with the overall treatment duration of orthognathic surgery? (Yes or No)
5. Which one is more important for you?
  - a. facial changes
  - b. dental changes
  - c. both
  - d. none of the above

Figure 1: Questionnaire

Patients stayed in the hospital for 5 days after surgery. Rigid fixation and an interocclusal splint were applied for 2 weeks, and patients wore light elastics for 2 weeks. Exclusion criteria were patients with mature cleft lip and palate with craniofacial syndromes and patients with orthognathic surgery without orthodontic treatment or with distractor devices. Institutional ethical approval was granted for this work by Kalinga Institute of Medical Sciences.

## Statistical analysis

All the questionnaires responses were coded and then analyzed using SPSS version 20.0 (Version 20.0. Armonk, NY: IBM Corp). The significant values were taken at 95% confidence interval with a  $P < 0.05$  value. Chi-square test was used to analyze the difference between the conventional and SFA mode of treatment.

## Results

Nineteen patients gave their consent to participate in the study. Eleven had undergone conventional orthognathic surgery, and 8 had opted for SFA. The intention to undergo surgery only for esthetics was noted in 57.89% of the patients; only improvement of chewing function in 10.52% and both in 31.57% [Table 1 and Figure 4].

It was also noticed that all patients wanted to see immediate changes in their facial appearance and less treatment time to get the desired results.

When asked about the postoperative results; among the patients who had undergone conventional surgery, 18.18% saw esthetic improvement, 27.27% felt an improvement in their masticatory function, 18.18%

6. Do you want immediate changes in facial appearance? (Yes or No)
7. How do you evaluate your postoperative overall result in immediate surgery / 6-8 months later surgery? Please mark one of the four given answers:
  - a. Aesthetic improvement only
  - b. improvement of chewing/masticatory function only
  - c. both
  - d. none/don't know
8. How do you feel exactly about the surgical outcome after immediate surgical treatment/surgical outcome after 6-8 months? Please mark one grade of the scale from 0 (poor) to 10 (excellent).
 

0-1-2-3-4-5-6-7-8-9-10
9. How do your family & friends feel about the surgical outcome of your treatment after immediate surgery/6-8 months post surgery? Please mark one grade of the scale from 0 (poor) to 10 (excellent)
 

0-1-2-3-4-5-6-7-8-9-10
10. How would you assess your facial appearance before surgery? Please mark one grade of the scale from 0 (poor) to 10 (excellent).
 

0-1-2-3-4-5-6-7-8-9-10
11. How would you assess your facial esthetics after immediate surgical treatment/surgical outcome after 6-8 months? Please mark one grade of the scale from 0 (poor) to 10 (excellent).

Figure 2: Questionnaire

felt an improvement in both, and 36.36% were unsure. [Table 2 and Figure 5].

Among patients who had undergone for SFA; 62.5% saw esthetic improvement, 12.5% felt an improvement in their masticatory function, and 25.0% felt an improvement in both [Table 2 and Figure 5].

There was a significant difference seen in the patient's perspective regarding pre- and post-surgical results in the case of patients who opted for SFA. The before and after results for patients undergone SFA was more appreciated [Table 3 and Figure 6].

When the outcome of both the surgeries was compared on an individual level and on peer level a significant value of *P* was obtained, suggesting a better response in case of SFA individuals [Tables 4, 5 and Figures 7, 8].

When the patients were asked about their comfortness after the surgery and on taking photographs. The patients who opted were more comfortable both immediately after the surgery and on taking photographs [Tables 6, 7 and Figures 9, 10].

When the patients were assessed on the satisfaction of the overall treatment effects, time and procedures, 81.81% of

patients who had undergone conventional surgery were satisfied, and 63.63% recommended it. While 100% of the patients who had undergone SFA were satisfied and recommended it [Tables 8, 9 and Figures 11, 12].

Most of the patients in further remarks wrote about their confidence boost and increased self-esteem after the surgery. However, there were also a few who wrote about the disheartening experience during the decompensation period of the conventional treatment. One patient who had undergone SFA wrote: "Right after the surgery, I met my new self." It is the immediate effect of SFA that impresses the patient and help them to cooperate more.

### Discussion

The definition of real success in our field first was initiated from the recognition of the problem and strong will power to overcome such problems by finding solutions. In this perspective, the SFA was successful in terms of focusing on the patients' chief problem.<sup>[14]</sup> "Accountability" which is defined as the orthodontist's ability to explain clearly the specific reason and course of treatment in a predictable way is yet another factor in our field. Orthognathic surgery has a certain amount of relapse, instability, and unpredictability.

0-1-2-3-4-5-6-7-8-9-10

12. Are you satisfied with the effects of 'surgery first'/conventional orthognathic surgery? (yes or no)

13. Do you feel comfortable post immediate surgery /after 6-8 months post surgery to eat outside in public places, smile meeting others, take photographs and not cover your face?  
0-1-2-3-4-5-6-7-8-9-10

14. How much you feel confident after surgery?  
0-1-2-3-4-5-6-7-8-9-10

15. Assuming that orthognathic surgery is indicated, would you recommend it to others?  
(Yes or no.)

16. If you recommending orthognathic surgery, then which one you would recommend to others?  
a- conventional orthognathic surgery b- surgery first approach

17. Do you have further remarks concerning the orthognathic surgery service?

Figure 3: Questionnaire




Figure 5: Postoperative overall result

**Table 1: Primary Intention of Surgery**

Primary Intention	Type of Surgery		Total	<i>P</i>
	Conventional	SFA		
Aesthetic Improvement	6	5	11	0.425
Mastication	1	1	2	
Both	4	2	6	
Total	11	8	19	




Figure 4: Primary intention of surgery




Figure 6: Appearance of face after surgery (visual analog scale)

**Table 2: Postoperative Overall Result**

Type of Surgery	Aesthetic Improvement	Masticatory Improvement	Both	Don't Know	Total	P
Conventional	2	3	2	4	11	0.114
SFA	5	1	2	0	8	
Total	7	4	4	4	19	




Figure 7: Outcome of surgery (visual analog scale) on individual level




Figure 8: Outcome of surgery (visual analog scale) on peer review




Figure 9: Comfortness postsurgery




Figure 10: Confidence after surgery on taking photographs




Figure 11: Patient's satisfaction with the effects




Figure 12: Recommendation of the treatment after surgery

Almost all patients are satisfied with the surgical outcomes of orthognathic surgery. This fundamental dilemma concerning “accountability” as a factor can improve our practice and can make a new major progress in our field. In the case of the conventional orthognathic treatment procedure, SFA needs to show scientific evidence as to whether this procedure can have a definitive advantage in terms of accountability.<sup>[15-17]</sup> SFA is thus believed to be a paradigm shift in the world of orthognathic surgery.

Dental occlusion and facial esthetics can show immediate effects postsurgery when using an SFA.<sup>[17,18]</sup> The SFA

offers an alternative to the orthodontics-first approach for correction of Dentofacial deformities. The advent of postoperatively accelerated orthodontic tooth movement also reduces the difficulties and complications associated with and the time spent on postoperative orthodontics. Both the orthodontist and the maxillofacial surgeon using an SFA should be well-experienced and should cooperate closely with each other to achieve predictable and satisfactory outcomes.<sup>[19-21]</sup> Orthodontists should be aware

**Table 3 : Appearance of Face after the Surgery (VAS)**

Score	Frequency			P
	Conventional	SFA	Total	
6	1	0	1	0.00
7	2	2	4	
8	3	2	5	
9	3	3	6	
	Median score=7	Median score=8.5	Median score=8	

**Table 4: Outcome of the Surgery (VAS) on individual level**

Satisfaction Score	Frequency			P
	Conventional	SFA	Total	
6	3	0	3	0.01
7	6	1	7	
8	2	3	5	
9	0	3	3	
10	0	1	1	
	Median score=7	Median score=8.5	Median score=7	

**Table 5: Outcome of the Surgery (VAS) on peer views**

Satisfaction Score	Frequency			P
	Conventional	SFA	Total	
6	5	0	3	0.00
7	4	1	5	
8	2	3	5	
9	0	3	3	
10	0	1	1	
	Median score=7	Median score=8.5	Median score=7	

**Table 6: Comfortness Post-Surgery**

Satisfaction Score	Frequency			P
	Conventional	SFA	Total	
5	2	0	2	0.00
6	3	0	3	
7	3	0	3	
8	3	1	4	
9	1	5	6	
10	0	2	2	
	Median score=7	Median score=8.5	Median score=8	

of the orthognathic principles, limitations in orthodontic movement and plan postoperative orthodontic treatment to include alignment, incisor decompensation, arch coordination, and occlusal interdigitation. The maxillofacial surgeon should be experienced enough of performing designated osteotomy and intermaxillary fixation with occlusion bite plate on malaligned dental arches and providing the stability after skeletal reposition.

**Table 7: Confidence after Surgery on taking Photographs**

Satisfaction Score	Frequency			P
	Conventional	SFA	Total	
6	3	0	3	0.00
7	3	0	3	
8	4	1	5	
9	1	1	2	
10	0	6	6	
	Median score=7	Median score=10	Median score=8	

**Table 8: Patient's Satisfaction with the Effects**

	Type of Surgery			P
	Conventional	SFA	Total	
Yes	9	8	17	0.485
No	2	0	2	
	11	8	19	

**Table 9: Recommendation of the Treatment by the Patients**

	Type of Surgery			P
	Conventional	SFA	Total	
Yes	7	8	15	0.103
No	4	0	4	
	11	8	19	

The reduction of the overall orthodontic treatment period and immediate facial changes would be the most significant advantages of this approach. Dentofacial deformity affects patient quality of life in our society.<sup>[22,23]</sup> The high satisfaction rate according to the questionnaire given to the patients was ranged between 70% and 87%. There was a marked improvement in self-confidence in 67.5% of patients as a result of an improved appearance and chewing function. Patients responded that immediate surgical treatment had a great impact on their friends and family.<sup>[24,25]</sup> Satisfaction of those correlated significantly with patients' satisfaction.<sup>[26]</sup> From the questions asked we get to know that patients' mental attitude about his/her postoperative appearance may be influenced by the support from family postsurgery. In contrast, the prevalence of postoperative complications such as numbness, prickle-sensations, pain in the temporomandibular joint area, or restrictions in mouth-opening had no overall effect on satisfaction.<sup>[27]</sup>

The primary factor of satisfaction with surgery first was whether or not the result was perceived to be an esthetic improvement. If there was an esthetic improvement in facial features immediately, the satisfaction was high in comparison to the conventional orthognathic surgery, regardless of functional problems.<sup>[28,29]</sup> An important result of this study was that a correlation between age, sex, and response to the questionnaire was not found. Young male

patients wanted functional improvement, and older male patients showed no alteration in self-esteem or depression with surgery first technique. Young females desired for improvement in self-confidence and older female patients displayed improved self-esteem and diminished depressive symptoms due to surgical intervention.

It is well-known that satisfaction and the perception of surgical outcome depend on two important factors, preoperative expectations and the degree to which the procedure is explained by the orthodontist or the maxillofacial surgeon. Patients who are psychologically depressed before surgery tend to report a higher recovery burden and experience more difficulty with symptoms with social/self-concerns in the initial months of the postsurgery. Various factors may change postoperative satisfaction of the patients like written data about the sequelae and the recovery period.<sup>[30]</sup>

During conversations, the previously operated patients talked about their experiences, which helped on the emotional preparation of the patients about to undergo orthognathic surgery. This activity did not create negative impressions in the patients because it did not cause distress or anxiety. Visualizing treatment simulation before surgery did not affect the perception of symptoms or satisfaction postsurgery. After surgery first approach, most patients report an improvement in self-confidence, body and facial image, and social adjustment, but some patients are only temporarily satisfied and become litigious or violent.<sup>[31,32]</sup> Patients who receive an inadequate explanation of the surgery are prone to be emotionally unprepared and anxious after surgery, but orthognathic surgery may not be beneficial for patients who assume that it will solve most of their problems.

A similar study was conducted by Huang *et al.*,<sup>[33]</sup> in which there was a comparison of oral health-related quality of life and satisfaction in SFA and conventional orthognathic surgery patients. The quality of life in orthodontic-first group was deteriorated before orthognathic surgery whereas, in surgery-first group, the quality of life was immediately improved which lead to better satisfaction.

The limitations of the study include convenience sampling and a nonvalidated questionnaire thus, making the findings of this study exclusive to a particular sector of population. The result may not have a quantitative accurateness, but qualitatively, it will remain the same as this study is basically the comparison of the SFA and Conventional orthognathic surgery on patient perceptiveness.

## Conclusion

The most distinctive factors for patient satisfaction after orthognathic surgery and conventional orthognathic surgery were facial esthetics. The patient satisfaction was more in orthognathic surgery than conventional orthognathic surgery. The timing of treatment and immediate results

are important factors toward patient satisfaction. Function, esthetics and even psychological aspects should be considered equally when planning surgery.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## References

1. Cadogan J, Bennun I. Face value: An exploration of the psychological impact of orthognathic surgery. *Br J Oral Maxillofac Surg* 2011;49:376-80.
2. Esperão PT, de Oliveira BH, de Oliveira Almeida MA, Kiyak HA, Miguel JA. Oral health-related quality of life in orthognathic surgery patients. *Am J Orthod Dentofacial Orthop* 2010;137:790-5.
3. Alanko OM, Svedström-Oristo AL, Tuomisto MT. Patients' perceptions of orthognathic treatment, well-being, and psychological or psychiatric status: A systematic review. *Acta Odontol Scand* 2010;68:249-60.
4. Nagasaka H, Sugawara J, Kawamura H, Nanda R. "Surgery first" skeletal class III correction using the Skeletal Anchorage System. *J Clin Orthod* 2009;43:97-105.
5. Park KR, Kim SY, Park HS, Jung YS. Surgery-first approach on patients with temporomandibular joint disease by intraoral vertical ramus osteotomy. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2013;116:e429-36.
6. Lee S, McGrath C, Samman N. Impact of orthognathic surgery on quality of life. *J Oral Maxillofac Surg* 2008;66:1194-9.
7. Alves e Silva AC, Carvalho RA, Santos Tde S, Rocha NS, Gomes AC, de Oliveira e Silva ED. Evaluation of life quality of patients submitted to orthognathic surgery. *Dental Press J Orthod* 2013;18:107-14.
8. Rusanen J, Lahti S, Tolvanen M, Pirttiniemi P. Quality of life in patients with severe malocclusion before treatment. *Eur J Orthod* 2010;32:43-8.
9. Laufer D, Glick D, Gutman D, Sharon A. Patient motivation and response to surgical correction of prognathism. *Oral Surg Oral Med Oral Pathol* 1976;41:309-13.
10. Choi WS, Lee S, McGrath C, Samman N. Change in quality of life after combined orthodontic-surgical treatment of dentofacial deformities. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2010;109:46-51.
11. Kiyak AH, Bell R. Psychosocial considerations in surgery and orthodontics. In: Proffit WR, White RP, editors. *Surgical-Orthodontic Treatment*. St. Louis: Mosby; 1991. p. 421-37.
12. Luther F, Morris DO, Hart C. Orthodontic preparation for orthognathic surgery: How long does it take and why? A retrospective study. *Br J Oral Maxillofac Surg* 2003;41:401-6.
13. Khadka A, Liu Y, Li J, Zhu S, Luo E, Feng G, *et al.* Changes in quality of life after orthognathic surgery: A comparison based on the involvement of the occlusion. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011;112:719-25.
14. Sabri R. Orthodontic objectives in orthognathic surgery: State of the art today. *World J Orthod* 2006;7:177-91.
15. Hernández-Alfaro F, Guijarro-Martínez R, Molina-Coral A, Badia-Escriche C. "Surgery first" in bimaxillary orthognathic surgery. *J Oral Maxillofac Surg* 2011;69:e201-7.
16. Behrman SJ, Behrman DA. Oral surgeons' considerations

- in surgical orthodontic treatment. *Dent Clin North Am* 1988;32:481-507.
17. Proffit WR, Miguel JA. The duration and sequencing of surgical-orthodontic treatment. *Int J Adult Orthodon Orthognath Surg* 1995;10:35-42.
  18. Proffit WR, White RP Jr. Who needs surgical-orthodontic treatment? *Int J Adult Orthodon Orthognath Surg* 1990;5:81-9.
  19. Gava EC, Miguel JA, de Araújo AM, de Oliveira BH. Psychometric properties of the Brazilian version of the Orthognathic Quality of Life Questionnaire. *J Oral Maxillofac Surg* 2013;71:1762.e1-8.
  20. Liu Z, McGrath C, Hägg U. The impact of malocclusion/orthodontic treatment need on the quality of life. A systematic review. *Angle Orthod* 2009;79:585-91.
  21. Barbosa Tde S, Tureli MC, Nobre-dos-Santos M, Puppim-Rontani RM, Gavião MB. The relationship between oral conditions, masticatory performance and oral health-related quality of life in children. *Arch Oral Biol* 2013;58:1070-7.
  22. Murphy C, Kearns G, Sleeman D, Cronin M, Allen PF. The clinical relevance of orthognathic surgery on quality of life. *Int J Oral Maxillofac Surg* 2011;40:926-30.
  23. Ballon A, Laudemann K, Sader R, Landes CA. Patients' preoperative expectations and postoperative satisfaction of dysgnathic patients operated on with resorbable osteosyntheses. *J Craniofac Surg* 2011;22:730-4.
  24. Rustemeyer J, Martin A, Gregersen J. Changes in quality of life and their relation to cephalometric changes in orthognathic surgery patients. *Angle Orthod* 2012;82:235-41.
  25. Kavin T, Jagadesan AG, Venkataraman SS. Changes in quality of life and impact on patients' perception of esthetics after orthognathic surgery. *J Pharm Bioallied Sci* 2012;4 Suppl 2:S290-3.
  26. Feu D, Miguel JA, Celeste RK, Oliveira BH. Effect of orthodontic treatment on oral health-related quality of life. *Angle Orthod* 2013;83:892-8.
  27. Rustemeyer J, Gregersen J. Quality of Life in orthognathic surgery patients: Post-surgical improvements in aesthetics and self-confidence. *J Craniomaxillofac Surg* 2012;40:400-4.
  28. Soh CL, Narayanan V. Quality of life assessment in patients with dentofacial deformity undergoing orthognathic surgery – A systematic review. *Int J Oral Maxillofac Surg* 2013;42:974-80.
  29. Rivera SM, Hatch JP, Rugh JD. Psychosocial factors associated with orthodontic and orthognathic surgical treatment. *Semin Orthod* 2000;6:259-69.
  30. Hatch JP, Rugh JD, Clark GM, Keeling SD, Tiner BD, Bays RA. Health-related quality of life following orthognathic surgery. *Int J Adult Orthodon Orthognath Surg* 1998;13:67-77.
  31. Kiyak HA, Hohl T, West RA, McNeill RW. Psychologic changes in orthognathic surgery patients: A 24-month follow up. *J Oral Maxillofac Surg* 1984;42:506-12.
  32. Motegi E, Hatch JP, Rugh JD, Yamaguchi H. Health-related quality of life and psychosocial function 5 years after orthognathic surgery. *Am J Orthod Dentofacial Orthop* 2003;124:138-43.
  33. Huang S, Chen W, Ni Z, Zhou Y. The changes of oral health-related quality of life and satisfaction after surgery-first orthognathic approach: A longitudinal prospective study. *Head Face Med* 2016;12:2.