

Original Article

Effects of mini-implant anchorage on mandibular rotation and esthetic outcomes in hyperdivergent skeletal Class II malocclusion treatment

Vo Thi Thuy Hong¹, Nguyen Thi Thu Phuong², Pham Thi Thu Hang³, Pham Anh Tuan⁴, Hoang Viet⁵, Tran Thi Kim Lien⁶

¹Department of Orthodontics, National Hospital of Odonto-Stomatology, ²Department of Orthodontics, School of Dentistry, Hanoi Medical University, ³Department of Odontology, 108 Military Central Hospital, ⁴Department of Mathematics and Information, Military Medical University, ⁵Department of Orthodontics and Pedodontics, Faculty of Dentistry, Van Lang University, ⁶Department of Odontology, Military Hospital 103, Ha Noi, Vietnam.



***Corresponding author:**


Tran Thi Kim Lien,
Department of Odontology,
Military Hospital, Ha Dong, Ha
Noi, Vietnam.

dr.lien103@gmail.com

Received: 29 April 2024
Accepted: 25 July 2024
Epub Ahead of Print: 05 October 2024
Published:

DOI
10.25259/APOS_92_2024

Quick Response Code:



ABSTRACT

Objectives: This research aimed to investigate the effects of mini-implants (MIs) on mandibular position and changes in lower face height, as well as their relationship with the position of the maxillary molars and incisors in the camouflage treatment of patients with hyperdivergent skeletal Class II malocclusion.

Material and Methods: This retrospective study included 27 patients aged 15–38 years. The hyperdivergent patients were diagnosed with skeletal Class II malocclusion and treated with tooth compensation, tooth extraction, posterior repositioning of the maxillary incisors, and anchoring with MIs. Cephalometric analyses of these patients were performed before (T1) and after (T2) treatment. Measurements were taken at each treatment stage and analyzed.

Results: Significant correlations were observed in the mandibular position and lower face height ($P < 0.05$). The GoGnSN angle decreased ($-1.05 \pm 1.61^\circ$), the ANS-Me distance reduced (-1.98 ± 4.76 mm), and the NBaPtGn angle increased ($1.67 \pm 4.08^\circ$). In addition, the vertical height position of the maxillary first molar significantly decreased (6M-FH, -1.61 ± 3.74 mm), while the vertical height position of the maxillary incisor increased (U1-FH, 1.48 ± 2.11 mm).

Conclusion: During the camouflaging treatment of patients with hyperdivergent skeletal Class II malocclusion using MIs anchorage, MIs helped intrude the maxillary first molar and extrude the maxillary incisor, leading to a difference in a mandibular counterclockwise rotation. Controlling the rotation of the lower jawbone helps create an improved esthetic for the patient after treatment.

Keywords: Skeletal Class II malocclusion, Hyperdivergent, Extraction premolar, Mini-implant

INTRODUCTION

Malocclusion, a common dental condition, refers to the misalignment of teeth and improper fitting of upper and lower teeth. Among its various classifications, skeletal Class II malocclusion is particularly notable, characterized by a prominent upper jaw or a receding lower jaw. Patients with skeletal Class II malocclusion often present with hyperdivergent facial types. Characteristics of hyperdivergent skeletal Class II malocclusion include the protrusion of the upper jaw and upper lip, incompetent lip growth, poor chin morphology, small nasolabial angles, and an overall long face, which are often the primary concerns for patients.^[1,2]

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, transform, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

©2024 Published by Scientific Scholar on behalf of APOS Trends in Orthodontics

Hyperdivergent skeletal Class II malocclusion has always been challenging to treat in orthodontics due to its characteristics involving both sagittal and vertical dimensions.^[3] The key to correcting these inconsistencies is to induce counterclockwise rotation of the mandible, thereby improving the patient's profile.^[4] Conventional treatment of skeletal Class II malocclusion in hyperdivergent patients has primarily involved the extraction of premolars in alleviating dental crowding and improving occlusal relationships. This method creates space, enabling the retraction of anterior teeth and achieving better alignment; however, premolar extraction alone often fails to address the vertical component of the malocclusion adequately.

The conventional approach is often combined with the use of elastics or headgear to manage vertical dimensions, but these methods have limitations.^[5] Elastics depend heavily on patient adherence and can introduce unwanted side effects, such as tipping of teeth or increased vertical dimensions. Headgear, while effective in controlling vertical growth, suffers from low patient acceptance due to its bulky and conspicuous features.

In addition, orthodontic appliances such as braces or functional appliances are typically employed in conjunction with premolar extraction to guide tooth movement and improve jaw relationships.^[6] However, these appliances primarily influence the sagittal plane and are limited in their ability to exert vertical control. The reliance on patient compliance, particularly with removable appliances, further complicates achieving consistent and predictable results.

Mini-implants (MIs) have emerged as a revolutionary adjunct in orthodontic treatments. MIs transmit forces directly to the jawbone, maintaining the desired tooth position without causing excessive extrusion. Moreover, by providing stable anchorage points, MIs allow for precise tooth movements without relying on patient compliance. MIs have been frequently utilized to regulate the vertical height of the anterior and posterior teeth, thereby causing the mandibles to rotate in a counterclockwise direction and enhancing facial profiles.^[7] This technique holds promise for managing vertical facial dimensions in hyperdivergent patients, particularly when combined with premolar extraction.

This research aimed to investigate the effects of MIs on mandibular position and changes in lower face height, as well as their relationship with the position of the maxillary molars and incisors in the camouflage treatment of patients with hyperdivergent skeletal Class II malocclusion. Through a comprehensive analysis, this research can contribute to the optimization of therapeutic strategies, thereby improving the quality of care for patients with complex malocclusions.

MATERIAL AND METHODS

Study design and ethics

This is a retrospective study approved by the Institutional Review Board of 108 Hospital (IRB: 1798/QD-BV). Patients were recruited from the Orthodontic Department at the National Hospital Odonto-Stomatology Hanoi from 2017 to 2021. All subjects provided written informed consent after being thoroughly informed about the study procedure and their rights. This study was conducted in accordance with the ethical principles of the Declaration of Helsinki.

Study subjects

Twenty-seven patients participated in the study, with the age ranging from 15 to 38 years. The sample size was calculated on clincalc.com. Based on a previous study by Costa *et al.*,^[8] anticipated means and standard deviation were entered, the detection power was set at 80%, and the alpha value was set at 5%. Inclusion criteria were (1) hyperdivergent skeletal Class II malocclusion (ANB >3.6°, GoGn-Sn >37°), (2) absence of systemic or chronic conditions, (3) no previous orthodontic treatment, and (4) satisfactory oral hygiene and periodontal health.

Procedures

The patients received camouflage treatment, which included premolar extraction, retraction of the maxillary incisors, and maximum anchorage using MIs.

Straight wire arch technology was employed to treat all study participants using pre-adjusted MBT appliances with a slot size of 0.022 × 0.028 inches. Treatment began with leveling and aligning using a 0.012-inch nickel-titanium archwire, followed using a 0.019 × 0.025-inch stainless steel (SS) archwire.

MIs (1.6 mm in diameter, 8 mm, and 10 mm in length, Jeil, Korea) were placed bilaterally in the alveolar bone between the maxillary first molar and second premolar. Anterior hooks positioned 2 mm distal to the lateral incisors were installed into the upper arch. Maxillary anterior teeth were retracted using 250 g of force applied through elastic traction from the MIs head to the anterior hooks.

All measurement items were recorded 3 times, and the average value was used as the final measurement result. Cephalometric points, planes was described and explained in [Figures 1-5]. Measurement errors were required to be <0.3° for angles and <0.4 mm for distances. If the error exceeded these thresholds, the reference points were re-evaluated, and the measurements were repeated until they met the required accuracy.

Statistical analysis

Data were expressed as the mean ± standard deviation. Statistically significant differences were analyzed using the paired *t*-test. All statistical analyses were performed using the Statistical Package for the Social Sciences (SPSS) version 22.0 (SPSS Inc., Chicago, IL, USA). The level of significance was set at *P* < 0.05.

RESULTS

There was no significant change observed in the SNA and SNB angles (*P* > 0.05), while the ANS-Me distance showed a significant decrease of -1.98 ± 4.76 mm (*P* < 0.05). The NBaPtGn angle increased significantly by $1.67^\circ \pm 4.08^\circ$ (*P* < 0.05), and the Wits appraisal decreased significantly by -0.74 ± 1.45 mm (*P* < 0.05). The GoGn-SN angle also decreased significantly by $-1.05^\circ \pm 1.61^\circ$ (*P* < 0.05) [Table 1].

The distance from the upper lip to the E-line (Ls-E) significantly decreased by -1.83 ± 2.16 mm (*P* < 0.05). Conversely, the distance from the lower lip to the E-line (Li-E) significantly increased by -2.61 ± 2.48 mm (*P* < 0.05). In addition, the distance from the upper lip to the Y-axis (Ls-axis *y*) decreased

by -4.1 ± 5.79 mm, and the distance from the lower lip to the Y-axis (Li-axis *y*) decreased by -4.52 ± 5.43 mm, both indicating a significant retraction of the lips (*P* < 0.05) [Table 2].

The analysis presented in [Table 3] revealed several key insights into the relationships between different skeletal measurements. The first line of [Table 3] showed that GoGnSN significantly influenced ANS-Me (*P* < 0.05), confirming the suitability of this model. The coefficient for GoGnSN is 0.388, which was statistically significant at the 5% level (*P* < 0.05), indicating that for every 1-unit increase in GoGnSN, there was a corresponding 0.388-unit increase in ANS-Me. The *R*² value for this model was 0.252, meaning that GoGnSN accounts for 25.2% of the variability in ANS-Me.

In contrast, the second line of [Table 3] indicated that GoGnSN did not significantly influence NBa-PtGn (*P* = 0.073), making this model unsuitable for explaining the relationship between these variables.

The third line of [Table 3] demonstrates that 6M-FH had a significant effect on GoGnSN (*P* > 0.05), validating the use of this model. The coefficient for 6M-FH was 0.842, statistically significant at the 5% level (*P* < 0.05), indicating that for each 1-unit increase in 6M-FH, GoGnSN increases by 0.842 units.

Table 1: The cephalometric skeletal variables at T1 and T2 for the hyperdivergent skeletal Class II malocclusion patients (Mean±SD).

Group/Measurements	Hyperdivergent			P-values
	Pre-treatment (T1) (n=27)	Post-treatment (T2) (n=27)	(Post-treatment)-(Pre-treatment) (T2-T1)	
SNA (°)	80.06±14.59	81.83±2.8	1.78±14.34	0.525
SNB (°)	75.94±2.82	76.11±2.88	-0.17±1.47	0.552
ANB (°)	6.87±1.92	5.72±1.44	-1.15±1.28	0.000
NBa-PtGn (°)	81.39±3.36	83.06±3.84	1.67±4.08	0.044
GoGnSN (°)	40.22±2.24	39.17±2.92	-1.05±1.61	0.002
MP-FH (°)	29.02±3.47	27.93±3.83	-1.09±3.3	0.097
PP-MP (°)	31.26±3.11	30.74±3.61	-0.52±2.54	0.299
ANS-Me (mm)	65.33±6.91	63.35±4.5	-1.98±4.76	0.04
Wits	2.65±1.69	1.91±1.37	-0.74±1.45	0.014

SNA: Sella nasion point A, SNB: Sella nasion point B, ANB: A point, nasion, B point, NBa-PtGn: Nasion, Basion - Pterygoid, Gnathion, GoGnSN: Gothion, Gnathion, Sella, Nasion, ANS-Me: ANS point.

Table 2: The Cephalometric soft tissue variables at T1 and T2 for the hyperdivergent skeletal Class II malocclusion patients.

Group/Measurements	Hyperdivergent			P-value
	Pre-treatment (T1) (n=27)	Post-treatment (T2) (n=27)	(Post-treatment)-(Pre-treatment) (T2-T1)	
Nasolabial angle (°)	91.48±12.57	100.2±8.11	8.73±10.87	0.000
Ls-E (mm)	1.77±1.74	-0.06±1.5	-1.83±2.16	0.001
Li-E (mm)	4.35±2.11	1.74±1.67	-2.61±2.48	0.000
Ls- axis-Y (mm)	75.13±7.04	71.04±4.23	-4.1±5.79	0.000
Li- axis-Y (mm)	71.8±7.12	67.28±4.24	-4.52±5.43	0.000
Pog'- axis-Y (mm)	61.04±6.69	58.24±5.22	-2.8±4.53	0.004

Ls-E: Ls to E line, Li-E: Li to E line, E line: a line connects the nose tip to the chin

Table 3: The index of predictable equation correlates between skeletal and soft tissue.

Ending variable	Predictable Equation			
	R-square	Coefficients regression (a)	Constant (b)	P
ANS-Me	0.252	GoGnSN 0.388***	23.72***	0.000
NBa-PtGn	0.047	GoGnSN 0.155* (0.073)	68.84***	0.073
GoGnSN	0.186	6M-FH 0.842***	62.813***	0.000

Predictable equation: $Y=ax+b$. Y: Ending variable (ANS-Me, NBa-PtGn, GoGnSN), a: Coefficients regression, x: Predictor variable (GoGnSN, U1-SN,6M-FH), b: Constant, ANS-Me: ANS point to Menton, NBa-PtGn: Nasion Basion - Pterygoid Gnathion, GoGnSN: Gothion Gnathion Sella Nasion. * $p<0.05$; *** $p<0.001$.




Figure 1: Simulation image of a point on the bone on cephalometric radiographies. S: Sella, N: Nasion, Po: Porion, Co: Condylion, Ba: Basion, Go: Gothion, PNS: PNS Point, Or: Orbital, ANS: ANS point, A: A point, B: B point, Pog: Pogonion, Gn: Gnathion, Me: Menton.




Figure 3: Simulation image of soft-tissue profile on cephalometric radiographies. Sn: Subnasale, Ls: Labrale superius, Li: Labrale inferius, Pog': Pogonion'




Figure 2: Simulated image of dental index points on cephalometric radiographies. U1: Upper central incisor, L1: Lower incisor, Ls: Labrale superius, Li: Labrale Inferius, Ms: Molar superior, Mi: Molar Inferius.




Figure 4: The Y-axis is defined as perpendicular to the X-axis (forming a 7° angle with the Sella nasion [SN] line downward).

The R^2 value for this relationship was 0.186, meaning that 6M-FH explains 18.6% of the variance in GoGnSN.

DISCUSSION

Hyperdivergent skeletal Class II malocclusion is particularly challenging to treat in orthodontics due to its sagittal and vertical inconsistencies. These cases often involve skeletal Class II malocclusion with an extremely high mandibular plane angle (GoGnSN >37°), possibly a retrusive mandible, anterior open bite, and severe overjet.

In this study, MI anchorage was used to retract the anterior teeth, extrude the maxillary anterior segments, and intrude the mandibular posterior segments. The results showed that the 6M-FH distance reduced by -1.61 ± 3.74 mm post-treatment, indicating that the first maxillary molar's distance to the Frankfort plane decreased, and the molar intruded ($P < 0.05$). The U1-FH distance increased by 1.48 ± 2.11 mm,




Figure 5: Simulated image of the anterior-posterior dimensions of the skeletal structure on cephalometric radiographies. N: Nasion, A: A point, B: B point.

indicating the first maxillary incisor's distance to the Frankfort plane increased, and the incisors extruded ($P < 0.05$). Moreover, the GoGnSN angle reduced by $-1.05 \pm 1.61^\circ$, the ANS-Me distance decreased by -1.98 ± 4.76 mm, and the NBaPtGn angle increased by $1.67 \pm 4.08^\circ$ [Table 1]. These changes suggested the mandible rotated in a counterclockwise direction and the lower face height decreased. The treatment effectively improved the profile both vertically and sagittally.

In treating skeletal Class II malocclusions for patients with a retrognathic mandible and a convex facial profile, orthodontic camouflage can be performed with or without extractions using Class II elastics to coordinate the relationship between the maxilla and mandible.^[9] However, Class II elastics and conventional anchorage can generate side effects, including the loss of mandibular anchorage and extrusion of the anterior and posterior teeth. Nanda *et al.*^[10] reported that after 20 years, skeletal anchorage has proven effective in treating open bites and controlling vertical face height, particularly by targeting vertical control of the posterior segments.

Patients aged 15–38 years (mean age 21.7 years) were chosen for this investigation to lessen the impacts of growth. In addition, the upper jaw in the protrusion position was included, and the SNA angle was limited to $80.06 \pm 14.59^\circ$. Patients with hyperdivergence were included in this study, and the size of the mandibular plane angle was balanced to prevent the measurement results from being affected by differences in mandibular morphologic traits.

Patients with hyperdivergent skeletal Class II malocclusion are typically present with a backwardly rotated mandible rather than maxillary prognathism.^[11] In this study, a small but statistically significant level of intrusion was observed for the maxillary incisors (0.1 ± 3.84 mm). The estimated center of resistance for the six anterior teeth is located between the centers of resistance of the four incisors and canines.^[12]

Table 4: The Cephalometric teeth variables at T1 and T2 for the hyperdivergent skeletal Class II malocclusion patients.

Group/Measurements	Hyperdivergent			P-value
	Pre-treatment (T1) (n=27)	Post-treatment (T2) (n=27)	(Post-treatment)-(Pre-treatment) (T2-T1)	
U1-SN(°)	109.35±8.53	102.41±6.95	-6.94±8.5	0.000
U1-ANSPNS (°)	118.93±6.62	110.48±8.61	-8.44±8.38	0.000
L1-MP(°)	97.07±6.59	91.56±4.72	-5.52±5.88	0.000
U1-L1(°)	112.3±10.41	126.5±7.9	14.2±11.92	0.000
Is- axis-Y (mm)	64.9±6.85	59.06±3.84	-5.89±5.43	0.000
li - axis-Y (mm)	60.08±6.83	56.04±3.6	-4.79±5.09	0.000
Ms- axis-Y (mm)	37.88±5.11	37.98±3.84	0.1±3.84	0.894
Mi- axis-Y (mm)	37.96±5.64	37.91±5.71	-0.06±5.1	0.955
Ms-FH (mm)	44.94±5.56	43.33±3.54	-1.61±3.74	0.034
U1-FH (mm)	49.81±3.94	51.3±3.93	1.48±2.11	0.001

SN: Sella-nasion plane, MP: Mandibular plane, FH: Frankfort horizontal plane, U1: Upper central incisor, ANSPNS: ANS point to PNS point, L1: Lower incisor, Ls: Labrale superius, Li: Labrale Inferius, Ms: Molar superior, Mi: Molar Inferius, Ms point: Molar superior.

The applied force had two distinct components: A backward retractive force and an intrusive force. These forces caused bodily intrusion and retraction of the maxillary anterior teeth. From a biomechanical perspective, the types of dental movements, such as root-lingual tipping, crown-lingual tipping, or bodily translation, are ultimately determined by the relationship between the line of action of a force and the position of the center of resistance of a tooth. The occlusogingival position of MIs and the vertical height of the anterior hooks are important factors that can alter the force direction. For instance, using a short anterior hook increases the vertical component and reduces the horizontal component of the force, and vice versa. In this research, with the maxillary MIs located 8–9 mm apical to the bracket slot and the anterior hooks at 2 mm, the force line passed just under the center of resistance. This configuration caused bodily retraction of the anterior teeth with only slight linguoversion.

In this study, the maxillary first molar showed distal movement of 0.1 ± 3.84 mm [Table 4], which was statistically significant. Our findings suggest that the mandible rotated forward. The hinge structure of the temporomandibular complex can explain the relationship between the counterclockwise rotation of the mandible and the reduction in molar and incisor positions (6M-FH, U1-FH) [Table 4]. During the process of retraction using MIs, the intrusive movement of the molars increased the distance between the hinge and the terminal fulcrum, thereby regulating mandibular inclination through the wedge effect. These findings suggest that vertical control of the maxillary incisor and first molar has a fundamental effect on mandibular rotation. The results led to forward and upward movement of the chin, which contributed to the improvement in the facial profile.

Liu *et al.*^[13] reported that controlling vertical dimensions has always been very difficult in orthodontics. When using conventional anchorage, the extrusion of the maxillary teeth causes the mandible to rotate in a clockwise direction, adversely affecting facial profiles. In the en-masse retraction process, the force line between MIs and the hook intrudes and pulls back the anterior teeth, effectively controlling vertical dimension and improving facial profiles.

Lu *et al.*^[14] reported that MIs made the angle of the mandibular related to the skull plane decrease -1.15° oscillate from -1.67° to -0.63° . In comparison, GoGnSN reduced $-1.05 \pm 1.61^\circ$ [Table 1]. In addition, a study of 15 research papers involving 489 patients reported that the first maxillary molar intrusion averaged -1.45 mm, oscillating between -2.02 mm and -0.89 mm ($P < 0.001$). In our results, the first maxillary molar intrusion was -1.61 ± 3.74 mm with $P < 0.05$ [Table 4].

According to the predictive equation, when the GoGnSN angle increased by 1 unit, the ANS-Me distance increased

by 0.388 units, with a coefficient of $R^2 = 0.252$. This indicated the GoGnSN angle explained 25.2% of the changes in the ANS-Me distance. Similarly, when the 6M-FH distance increased by 1 unit, the GoGnSN angle increased by 0.842 units, with a coefficient of $R^2 = 0.186$, indicating that 6M-FH explained 18.6% of the changes in the GoGnSN angle. The GoGnSN angle's influence on the NBa-PtGn angle was not significant ($P = 0.073$), indicating this model was not suitable for predicting changes in NBa-PtGn [Table 3].

CONCLUSION

In the group with the open mandibular angle and anchored with MIs, the mandible was not rotated downward and backward, the maxillary incisors pulled back more, and the 6th teeth moved forward less than the traditional anchorage group. NBaPtGn facial axial angle decreased compared to before treatment, GoGnSN angle remained unchanged, and ANS-Me lower facial floor height remained the same.

Ethical approval

The research/study was approved by the Institutional Review Board at 108 Hospital, number 1798/Qu0110-BV, dated March 20, 2019.

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Use of artificial intelligence (AI)-assisted technology for manuscript preparation

The authors confirm that there was no use of artificial intelligence (AI)-assisted technology for assisting in the writing or editing of the manuscript and no images were manipulated using AI.

REFERENCES

1. Gkantidis N, Halazonetis DJ, Alexandropoulos E, Haralabakis NB. Treatment strategies for patients with hyperdivergent Class II Division 1 malocclusion: Is vertical dimension affected? *Am J Orthod Dentofacial Orthop* 2011;140:346-55.
2. Buschang PH, Jacob H, Carrillo R. The morphological

- characteristics, growth, and etiology of the hyperdivergent phenotype. Netherlands: Elsevier; 2013. p. 212-26.
- Mishra D, Natarajan M, Urala AS. Lip profile changes in patients with Class II Division 1 malocclusion of varied growth patterns treated with maxillary premolar extractions: A pilot study. *Am J Orthod Dentofacial Orthop* 2020;158:684-93.
 - Rask H, English JD, Colville C, Kasper FK, Gallerano R, Jacob HB. Cephalometric evaluation of changes in vertical dimension and molar position in adult non-extraction treatment with clear aligners and traditional fixed appliances. *Dental Press J Orthod* 2021;26:e2119360.
 - Fontes FP, Bellini-Pereira SA, Aliaga-Del-Castillo A, Patel MP, Freitas MR, Henriques JF, *et al.* Comparison of the dentoskeletal and soft tissue changes with the cervical headgear and Jones Jig followed by fixed appliances in Class II malocclusion patients: A retrospective study. *Int Orthod* 2020;18:424-35.
 - Männchen R, Serafin M, Fastuca R, Caprioglio A. Does early treatment improve clinical outcome of Class II patients? A retrospective study. *Children (Basel)* 2022;9:232.
 - Akbaydogan LC, Akin M. Cephalometric evaluation of intrusion of maxillary posterior teeth by miniscrews in the treatment of open bite. *Am J Orthod Dentofacial Orthop* 2022;161:621-7.
 - Costa JV, Ramos AL, Iwaki Filho L. Zygomatic-maxillary cortical bone thickness in hyper, normo and hypodivergent patients. *Dental Press J Orthod* 2021;26:e211965.
 - Janson G, Sathler R, Fernandes TM, Branco NC, de Freitas MR. Correction of Class II malocclusion with Class II elastics: A systematic review. *Am J Orthod Dentofacial Orthop* 2013;143:383-92.
 - Nanda R, Castroflorio T, Garino F, Ojima K. Principles and biomechanics of aligner treatment-e-book. Netherlands: Elsevier Health Sciences; 2021.
 - Plaza SP, Reimpell A, Silva J, Montoya D. Relationship between skeletal Class II and Class III malocclusions with vertical skeletal pattern. *Dental Press J Orthod* 2019;24:63-72.
 - Bouton A, Simon Y, Goussard F, Teresi L, Sansalone V. New finite element study protocol: Clinical simulation of orthodontic tooth movement. *Int Orthod* 2017;15:165-79.
 - Liu Y, Yang ZJ, Zhou J, Xiong P, Wang Q, Yang Y, *et al.* Comparison of anchorage efficiency of orthodontic mini-implant and conventional anchorage reinforcement in patients requiring maximum orthodontic anchorage: A systematic review and meta-analysis. *J Evid Based Dent Pract* 2020;20:101401.
 - Lu Y, Zhang W, Zhao B, Liu Y. Vertical control of a severe hyperdivergent skeletal Class II malocclusion with steep posterior occlusal plane in a camouflage case. *Medicina (Kaunas)* 2022;58:1217.

How to cite this article: Hong VT, Lien TT, Phuong NT, Hang PT, Tuan PA. Effects of mini-implant anchorage on mandibular rotation and esthetic outcomes in hyperdivergent skeletal Class II malocclusion treatment. *APOS Trends Orthod*. doi: 10.25259/APOS_92_2024